

## Influence of herbal adjuncts in shaping the quality and bioactive properties of craft beers

<sup>1</sup>Katarzyna Gościńska\* , <sup>1</sup>Paulina Bracka, <sup>2</sup>Katarzyna Mikołajczyk-Bator ,  
<sup>3</sup>Krunoslav Aladić 

<sup>1</sup>Department of Agronomy and Food Technology, Faculty of Agriculture and Biotechnology,  
Bydgoszcz University of Science and Technology, Kaliskiego 7, 85-796, Bydgoszcz, POLAND

<sup>2</sup>Department of Natural Science and Quality Assurance Poznań University of Economics and Business,  
Al. Niepodległości 10, 61-875, Poznań, POLAND

<sup>3</sup>Faculty of Food Technology Osijek, Josip Juraj Strossmayer University of Osijek, Franje Kuhaca 20, 31000 Osijek, CROATIA

\*Corresponding author: Katarzyna Gościńska, e-mail: katarzyna.goscińska@pbs.edu.pl

**Abstract.** The aim of this study was to conduct a comprehensive evaluation of selected herbal adjuncts on the quality profile of craft beer. The assessment included the determination of total polyphenol content (TPC), characterization of antioxidant activity and evaluation of the influence of herbal adjuncts on the sensory attributes of the final product. This research refers to innovative trends in craft brewing, oriented towards development of products with enhanced functional value and unique organoleptic profile, which is crucial both in the context of the expanding fermented beverage market and growing consumer interest in health-promoting foods. The base beer was Pale Ale (PA), enriched with plant-derived additives (thyme, ginger, lavender, marjoram, mint). The resulting beverages were tested for total extract and alcohol content, TPC, flavonoids, pH, antioxidant activity (FRAP, ABTS) and color expressed in EBC (European Brewery Convention) units. Sensory evaluation was carried out with 100 panelists using a five-point hedonic scale. Based on obtained results, it was found that the addition of thyme and ginger was the most beneficial in improving health-promoting properties of beers. In terms of sensory profile mint and ginger were the most favourable. The findings are important for the development of innovative technologies in craft brewing and also for the creation of products that meet the growing consumer interest in functional and health-promoting food. Despite the declared beneficial health effects, beer should be consumed in moderation due to the adverse effect of ethanol on the human body and the risk of alcohol dependence.

**Keywords:** craft beer, herbs, sensory evaluation, polyphenols, antioxidant activity

### INTRODUCTION

Beer is one of the most popular alcoholic beverages in the world, valued for its flavor and the presence of natural nutritional and biological compounds such as carbohydrates, amino acids, vitamins and polyphenols. Traditional beer is produced from water, barley malt, hops and yeast (Olšovska et al., 2014; Djordjevic et al., 2016; Di Domenico et al., 2020; Ullah et al., 2020). Nowadays the brewing

industry is undergoing dynamic changes. Craft beers are gaining increasing importance due to their flexible formulations and the potential for incorporating innovative ingredients.

According to reports by the Polish Office of Competition and Consumer Protection (UOKiK) and eLeader (using the Shelf Recognition AI app), craft beers are gaining popularity and occupying more shelf space in stores. However, their overall market share remains relatively small,

received 3 November 2025; reviewed 14 December 2025; accepted 22 December 2025

Authors declare no conflict of interest.



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

and the overall production of beer in Poland has seen a decline in recent years (UOKiK, eLeader, 2023). This trend reflects growing consumer interest in natural ingredients and awareness of the potential health effects of high-proof alcohol (Elzinga et al., 2015; Gomez-Corona et al., 2016; Preiss et al., 2018).

Modern consumer expects beer not only to be refreshing but also to offer a unique sensory profile (distinctive flavor and aroma), innovative ingredients and potential health benefits (Wojtyra, 2020; Brezinova, 2021; Kikut-Ligaj, Mikołajczyk-Bator, 2022; Mazengia et al., 2022). In consumers' opinion, craft beers are most often chosen because of their variety of flavors and superior quality compared to industrial beers (Kikut-Ligaj, Mikołajczyk-Bator, 2022).

The technological flexibility of small breweries enables producers to quickly respond to changing market demands, experiment with the formulations and develop products with increased functional value. The market offers beverages crafted from various types of malt and hops, as well as adjuncts including herbs, fruits, spices and bee products (Mastanjevic et al., 2019; Nardini, Garaguso 2020; Pluhackova et al., 2020). The additives used not only impart a characteristic flavor and aroma to the beers (Carvalho et al., 2016), but also enrich them with bioactive compounds such as polyphenols, flavonoids, vitamins with high antioxidant activity and enhance the shelf life of the products (Nardini, Garaguso, 2020; Horincar et al., 2020; Lin et al., 2023). Herbs, especially from *Lamiaceae* (L.) family represent a particularly interesting group of additives – due to their content of essential oils, polyphenols and other secondary metabolites with confirmed antioxidant, antibacterial, anti-inflammatory properties (Sohrabvandi et al., 2012; De Falco et al., 2013; Galovicova et al., 2021).

Research on the use of herbs and spices in craft brewing is of significant importance to the industry, as it not only allows for the expansion of the product range and its better adaptation to consumers' expectations, but also enables the creation of beers with increased functional value, which can provide competitive advantage in the rapidly growing market for fermented beverages. Moreover, incorporating plant ingredients into formulations can be seen as a synergistic effect between tradition and modernity – combining classic brewing technology with current health and nutritional trends. Following this trend of searching for an innovative craft beer recipes with an attractive sensory profile and increased functional value, the five Pale Ale (PA) variants were developed with the addition of plant ingredients: thyme, ginger, lavender, marjoram and mint.

## MATERIALS AND METHODS

### Materials

The subject of the study was beer with herbal adjuncts. PA beer was selected as a base beer. PA barley malt (Weyermann, Germany), water, Hercules hops and selected

brewing yeast strains *Saccharomyces cerevisiae* (SafAle US-05, Fermentis) were used in the production process of the beer. In the experimental variants selected lyophilized plant materials with health-promoting properties were added, including thyme (0.50 g/L), ginger (1.00 g/L), lavender (0.10 g/L), marjoram (0.30 g/L) and mint (0.50 g/L). The given values refer to the mass of the dry plant material per 1 liter of wort before fermentation process. The plant parts used were leaves for thyme, marjoram, and mint, rhizome for ginger, and flowers for lavender. After lyophilization, the plant material was ground into a coarse powder. The amounts of additives were determined based on preliminary studies involving a series of samples with varying proportions of plant material (0.1–2 g/L). The assessment of optimal doses was performed using sensory tests (N = 20 people, 5-point hedonic scale).

### Beer production process

Beer production process was carried out according to the standard technology described by Lehl (2017), using a Cobra CB3 device (COBRA, HOMEBREWING).

Mashing was carried out using the infusion method following the temperature profile: a 15-minute rest at 52 °C to activate proteolytic enzymes, followed by a 30-minute rest at 63 °C to hydrolyze starch into dextrins and fermentable sugars and heating to 72 °C for 20 minutes to promote  $\alpha$ -amylase activity. After reaching 78 °C, filtration and sparging were performed. The wort was boiled for 60 minutes with hops added in three equal doses of 20 g each, according to the brewing protocol. After boiling the wort was cooled down to 20 °C and aerated, then inoculated with a top-fermenting *Saccharomyces cerevisiae* yeast strain. Vigorous fermentation was carried out at 18–20 °C for 7 days, after which the beer was decanted from the sediment and transferred to secondary fermentation conducted at 12 °C for another 14 days. Lyophilized herbs (thyme (*Thymus vulgaris* L.), ginger (*Zingiber officinale* Rosc.), lavender (*Lavandula angustifolia* Mill.), marjoram (*Origanum majorama* L.), mint (*Mentha  $\times$  piperita* L.) were added directly to the fermenters during secondary fermentation, in doses according to the formulation. After maturation, the beer was filtered to remove suspended particles and any herb residues, then bottled with added 5% sucrose for re-fermentation. The finished product was conditioned at 4 °C for a minimum of 14 days.

### Physicochemical analysis

In the finished products, pH, real extract content, total polyphenol content (TPC), flavonoids content, antioxidant activity using the ABTS<sup>+</sup> cation radical and FRAP methods and as well as color in the EBC system were determined. In addition, the sensory evaluation was carried out.

The alcohol content in beer samples was determined according to the analytical methods EBC 9.43.2 (EBC An-

alytica, 2010) using Alcozyzer Beer with a DMA 4500M density module (Anton Paar GmbH, Austria). Real extract of wort was calculated based on the determined alcohol content in beer in compliance with EBC 9.43.2 method (EBC Analytic, 2010). Active acidity (pH) was measured according to the PN-90/A-75101/06 standard. The method is based on testing the potential difference between two electrodes immersed in the tested solution. The assessment was performed using an EDGE HI 2002 pH meter (Hanna Instruments, Italy).

Total polyphenol content was quantified using a method described by Fang et al. (2006) based on the reaction with a Folin-Ciocalteu reagent. The absorbance of the complex was measured at a wavelength of 735 nm, and the results were expressed in mg of chlorogenic acid per liter of beer (mg/L) using a calibration curve prepared for this compound. The flavonoid content was determined using the spectrophotometric method described by Papoutsis et al. (2016) using NaNO<sub>2</sub> and AlCl<sub>3</sub> solutions, measuring the absorbance at a wavelength of 510 nm. Beer samples were diluted 1:5 with distilled water before the analysis. The results were expressed as mg of quercetin equivalents (Sigma-Aldrich, St. Louis, MO, USA).

Antioxidant activity was expressed by two methods. FRAP method was carried out in accordance with the procedure described by Gościnnna et al. (2019), based on Benzie and Strain (1996), measuring absorbance at a wavelength of 593 nm and presenting the results in mmol Fe<sup>2+</sup>/L of beer using a calibration curve prepared for iron ions. For the ABTS<sup>•+</sup> cation radical assay, beer samples were first degassed by ultrasonic treatment and centrifuged at 4000 rpm for 15 min at 4 °C (Hettina Zentrifugen, Rotina 420 R, Germany). The obtained supernatant was used as the beer extract. The ABTS<sup>•+</sup> assay was then performed as described by Re et al. (1999) by incubating the ABTS<sup>•+</sup> working solution with the beer extract at 30 °C, followed by absorbance measurement at 734 nm. The results were expressed as  $\mu$ mol Trolox/L of beer.

All measurements were performed using a SHIMADZU UV-1800 spectrophotometer (UV-Vis Spectrophotometer System, Japan). Each analysis was conducted in triplicate for each sample.

### Sensory evaluation

Sensory analysis was conducted after the beers had completed secondary fermentation, were filtered, bottled with added sugar for re-fermentation, and conditioned for at least 14 days at 4 °C.

### Consumer hedonic assessment

Consumer acceptance was evaluated using a five-point hedonic scale (5 – very satisfactory, 4 – satisfactory, 3 – average, 2 – unsatisfactory, 1 – very unsatisfactory). The evaluation panel consisted of 100 individuals (women and

men aged 19–69) who were regular consumers of this type of product.

Each participant evaluated all four beer samples, including a control PA beer without herbal additives. Samples were served chilled in a randomized order. Portions of 30 mL were served in transparent, tulip-shaped glasses (100 mL capacity) marked with randomly generated two-digit codes.

### Descriptive sensory analysis

A descriptive sensory analysis was conducted to characterize the sensory attributes of the examined beers, including flavor, aroma, color, and foam characteristics. The evaluation was performed by a trained panel of 100 assessors experienced in beer sensory analysis.

Panelists evaluated the samples qualitatively using predefined sensory descriptors commonly applied in beer assessment (e.g., bitterness, herbal notes, spiciness, floral aroma, foam density and color).

Attributes were considered “not perceived negatively” when no undesirable or off-flavors were reported by the panel during the descriptive assessment.

### Statistical analysis

The influence of the analyzed factor on the physicochemical properties of products was determined using analysis of variance (ANOVA) in Statistica 13.1 software. Tukey's test was also applied with a significance level of  $\alpha = 0.05$  to examine the differences between mean values. Relationships between the examined qualitative characteristics were analyzed using Spearman rank correlation coefficient ( $p < 0.05$ ). The results were presented as the arithmetic means  $\pm$  standard deviation (SD) from 3 replicates for each sample. For sensory evaluation principal component analysis (PCA) were performed.

## RESULTS AND DISCUSSION

### Characteristics of the base PA beer

The physicochemical characteristics of laboratory produced PA beer are presented in Table 1, Figure 1A, B and 2. Real extract content was 3.23% and the alcohol content was 4.94% V/V. The pH was 4.37. The beer's color was 14.33 in EBC which corresponds to a dark golden, amber color (Table 2).

Obtained values of the extract and alcohol are within the typical range for PA beers, which indicates that the fermentation process proceeded correctly and that the brewing process was stable. The pH value of 4.37 is compliant with the expected range for this style of beer ensuring adequate microbiological stability and positively influencing the flavor and freshness of the beverage (Lehl, 2017; Cocuzza et al., 2019).

Table 1. The results of physicochemical measurements for tested beers with functional additives.

Beer variant	Real extract [%]	Alcohol [% V/V]	pH	Color (EBC)
PA*	3.23±0.04 a	4.94±0.10 b	4.37±0.03 cd	14.33±0.38 b
PAT	3.48±0.09 b	4.74±0.12 a	4.33±0.04 bc	15.37±0.98 ab
PAG	3.25±0.08 a	4.75±0.07 a	4.30±0.01 ab	16.63±1.06 c
PAL	3.29±0.09 a	4.61±0.05 a	4.33±0.02 bc	16.83±0.58 c
PAMR	3.34±0.08 ab	4.70±0.10 a	4.26±0.05 a	13.87±1.16 ab
PAM	3.33±0.14 ab	4.74±0.06 a	4.39±0.04 d	12.57±0.67 a

\*PA – Pale Ale (control), PAT – Pale Ale with thyme, PAG – Pale Ale with ginger, PAL – Pale Ale with lavender, PAMR – Pale Ale with marjoram, PAM – Pale Ale with mint. The values are means ± SD of three independent replicates. a, b, ... means sharing the same letter in column are not significantly different (p<0.05)

Table 2. Coloring of beers according to the EBC (Rum, 2005).

EBC	Color	Description	Beer example
1–4		Straw, light yellow, light gold	Pale lager
5–8		Yellow, gold	Pilsner
9–20		Dark gold, dark yellow, amber	Wheat
21–35		Light brown, brown, copper	Pale ale
36–60		Brown	Bock
> 60		Dark brown, black	Stout




Figure 1. (A) Total polyphenol content (TPC) and (B) flavonoids in analyzed beer variants. A, B, ..., a, b, ... – means sharing the same letter in column are not significantly different (p<0.05). Abbreviations – see Table 1.

The measured EBC color (14.33) is slightly lower than the typical range reported for Pale Ale (20–35 EBC, Table 2). This deviation may be attributed to the specific selection of malts and brewing conditions used in this laboratory-scale production, which produced a slightly lighter shade of amber. Nevertheless, the resulting color remains within the spectrum of craft beer variations, which can differ depending on the raw materials, adjuncts, and brewing methods employed (Ualema et al., 2024).

Total polyphenol content was 609.1 mg/L, and flavonoids content was 52.8 mg/L (Figure 1A, B). Antioxidant capacity measured using the ABTS<sup>•+</sup> cation radical was equal to 5.5 μmol of Trolox/L, while FRAP reducing capacity – 665.7 mmol Fe<sup>2+</sup>/L (Figure 2). The results indicate that the analyzed PA beer was characterized by a high content of bioactive compounds and strong antioxidant potential, which, when considered together with its color profile, are generally consistent with the characteristics of beers of this style. Although the measured EBC value was slightly lower than the typical range reported for Pale Ale, the observed color remains within the spectrum of variation commonly found in craft beers. These findings are consistent with the literature data, which indicate that PA beers contain increased content of polyphenolic compounds, that determine significant sensory profile of beer, including its color (Di Domenico et al., 2020; Habschied et al., 2021).

Many factors influence the polyphenolic content in beers, as well as the antioxidant capacity. The type and quality of malt used are of key importance. In the study by Ditrych et al. (2016) the lowest content of polyphenolic compounds, at 74 mg/L was observed in non-alcoholic beer, which was due to the low amount of malt used during production. In contrast the highest concentration of polyphenols was detected in dark beers, where the values ranged from 233.9 mg/L to 407.5 mg/L (Ditrych et al., 2016). The variety and quantity of hops used, as well as the conditions of the mashing process (temperature, time, pH), the method of wort filtration, the fermentation process and maturation, are also of significant importance (Sokol-Lętowska, 2013; Skendi et al., 2017; Ramos da Silva et al.,

2021). The higher proportion of specialty malts containing melanoids and reductones formed by Maillard reaction during additional thermal processing of malt (e.g. roasting) promotes an increase in polyphenol content (Shopska et al., 2021; Guido, Ferreira 2023). These compounds are responsible for the darkening of beer color (Coronas et al., 2024). However, some stages of the process, such as filtration and clarification may lead to their partial loss (Yağınçıray et al., 2022; Kumari, Sit, 2025). Parameters like temperature and pH during mashing, determine the extraction of phenolic compounds from the malt and hops, which influences the final biological activity of beer (Šibalić et al., 2021; Gribkova et al., 2022).

In this context it is worth pointing out the differences between industrial and craft beers. The craft beers are usually brewed in smaller batches and use less aggressive clarification methods, with higher use of specialty malts and more aromatic hops varieties. Industrially produced beer needs more intense stabilization and filtration methods, because of the desire for repeatability, stability and high clarity, which reduces the content of polyphenols, flavonoids and antioxidant activity of beverages (Humia et al., 2019). These factors do not favor maintaining a high level of bioactive compounds, as confirmed by scientific studies. As reported by Ambra et al. (2021) during industrial-scale brewing, approximately 60% of polyphenols from malt are degraded due to high-temperature pasteurization and filtration. Craft beers do not undergo these processes and maintain higher concentration of polyphenolic compounds. Sperkowska et al. (2021) studied 8 industrial beers and noticed significantly lower concentration of bioactive compounds than those reported in her own studies. The average polyphenol content in her studies was 237.2 μg GAE/L (gallic acid equivalents), while for flavonoids it was 45.03 μg CAT/L (catechin equivalent). The antioxidant capacity measured by FRAP method was 1.2 μmol Trolox/L.

Based on the results presented in Figures 1 and 2, it was observed that herbal adjuncts had positive influence on the polyphenol and flavonoids content and also on the antioxidant capacity of studied beers, in comparison to base beer. At the same time, a statistically significant decrease in alcohol content was found in the samples with plant additives relative to the base beer. The highest alcohol concentration was observed in beers containing ginger rhizoma (4.8% V/V), representing a 3.9% decrease compared to the base beer. Conversely, the lowest concentration was found in beers with lavender flowers (4.6% v/v), corresponding to a 6.7% decrease relative to the base beer.

These findings suggest that the presence of herbal adjuncts could have affected the fermentation process and sugar assimilation by yeast. As stated by Viejo et al. (2018) and Anderson et al. (2019), alcohol content is influenced not only by its concentration in the wort but also by the type of malt, technological process, and yeast strain used. Bottom-fermented beers (e.g., lager, pilsner, porter) typically have a higher concentration of residual sugars and

lower alcohol content (approx. 3.5–4.5% v/v) and are brewed with *Saccharomyces pastorianus*. In contrast, top fermented beers (e.g. ALE, IPA) are characterized by higher alcohol concentration (approx. 4–7% V/V) due to the activity of *Saccharomyces cerevisiae* yeast strain.

Each variant of beer with plant additives showed a slight increase in real extract content compared to the base beer (3.2%). The highest concentration was observed in beers with thyme (3.48%, increase by 7.2%) and the lowest content in the beers with ginger (3.25%, increase by 0.6%) (Table 1). The results suggest may indicate that herbal addition affects the extraction of soluble substances from raw materials and their availability during the fermentation. The thyme beer variant showed a higher real extract content (statistically significant differences) compared to the ginger (3.25%), lavender (3.29%) and control beers (3.23%) (Table 1).

The pH value for all beer types ranged from 4.26 to 4.39 (Table 1). All beers (except the one with mint addition, which showed a slight pH increase of 0.02) exhibited a small decrease in pH compared to the base beer. The largest decrease was observed in beers with marjoram (4.26; 0.11 lower) and ginger (4.30; 0.07 lower). Statistically significant differences were also noted between beers with marjoram and mint (4.29). According to Basarova et al. (2015), the pH value of beer primarily depends on organic acids produced during fermentation and typically falls within 4.0–4.9. Minor variations in pH were observed among beers with different herbal additions. The beer with marjoram (PAMR) showed the lowest pH (4.26), which was statistically lower than several other variants, while the remaining samples did not differ significantly from the control (PA, 4.37) (Table 1).

Herbal adjuncts caused distinct changes in the color of finished products (Table 1). Beers with thyme, lavender and ginger addition had a higher EBC value than base beer (15.37, 16.63, 16.83, 14.33 accordingly). Similar effect was observed in the studies of Gościńska et al. (2019) where the addition of thyme also resulted in the darkening of the beer. However, mint and marjoram addition resulted in lightening of beers (12.57 and 13.87 EBC accordingly). Statistically significant differences were observed between base beer and beers with ginger, lavender and mint and also amongst beer with marjoram or thyme and beer with ginger or lavender (Table 1). The differences in EBC values may be related to the presence of natural pigments in plant tissues and, as previously mentioned, to Maillard reaction and oxidation of polyphenols during the brewing process. The technological process also has a significant impact on color, as it can accelerate the oxidation of polyphenols and formation of complex compounds that influence the final color (Pieczonka et al., 2021).

Herbal adjuncts significantly affected the content of polyphenolic compounds compared to the base beer (609 mg/L) (Figure 1A). The highest polyphenol content was found in beer with thyme (1107.8 mg/L, an 80% increase), and

the lowest in beer with mint (905.2 mg/L, a 48% increase). Beers with thyme, marjoram, and ginger contained significantly more polyphenols than those with lavender or mint, while no significant differences were found between mint and lavender or between ginger and marjoram beers.

A similar trend was observed for flavonoids (Figure 1B). The base beer contained 52.8 mg/L, while beer with thyme had the highest concentration (72.9 mg/L, 38% increase), and beer with mint the lowest (60.9 mg/L, 15% increase). Statistical analysis revealed significant differences between the base beer and all beers with herbal adjuncts, as well as among individual additives, except for beers with ginger (71.1 mg/L) and thyme (72.8 mg/L), where no significant difference was observed.

Similar observations were made by Solgajova et al. (2024), who demonstrated a significant increase in polyphenol content after the addition of dried herbs (sage, oregano, lemon balm, thyme) during fermentation. Compared to the control beer (253.6 mg/L), this increase ranged from 118% for thyme (553.8 mg/L) to 216% for lemon balm (763 mg/L). However, Gościńska et al. (2019) reported that adding fresh thyme (7 g/L) increased TPC by only 3.5% compared to the base beer (321.2 mg/L). Djordjević et al. (2016) achieved a 37% increase in TPC after adding liquid thyme extract (0.5 mL/L). Such variations indicate that the quality, form, and processing of the additive (e.g., maceration time) are critical factors influencing the final effect.

Plant adjuncts can be added at any stage of brewing; however, studies indicate that the earlier they are added, the greater the potential loss of polyphenolic compounds due to oxidation and degradation. Adding plant material during mashing may therefore negatively affect the concentration of bioactive compounds and antioxidant activity (Gorjanovic et al., 2010; Sanna, Pretti, 2015).

In all samples, the addition of herbs to the base beer resulted in a statistically significant increase in antioxidant activity (ABTS<sup>+</sup>, FRAP) (Figure 2). The highest antioxidant activity was noted in beer with thyme (ABTS<sup>+</sup>: 7.16  $\mu\text{mol Trolox/L}$ , increase by 31%; FRAP: 918.3 mmol Fe<sup>2+</sup>/L, increase by 38%), while the lowest was found in beer with mint (ABTS<sup>+</sup>: 5.5  $\mu\text{mol Trolox/L}$ , increase by 0.7%; FRAP: 767 mmol Fe<sup>2+</sup>/L, increase by 15%) (Figure 2). For the FRAP method, significant differences in antioxidant capacity were observed for each beer variant. However, for the ABTS<sup>+</sup> method, no differences were noted between beers with ginger and marjoram or between those with lavender and mint. These results align with Lin et al. (2023), who reported that the addition of herbs such as jasmine and hibiscus enhances antioxidant activity, and Solgajova et al. (2024), who observed multiple increases in TEAC (ABTS) activity in Pilsner beers with added thyme, sage, oregano, or lemon balm.

Therefore, this research indicates that herbs are valuable brewing adjuncts due to their high content of bioactive sub-




Figure 2. Antioxidant activity FRAP and ABTS<sup>+</sup> cation radical of analyzed beer variants: PA – Pale Ale (control), PAT – Pale Ale with thyme, PAG – Pale Ale with ginger, PAL – Pale Ale with lavender, PAMR – Pale Ale with marjoram, PAM – Pale Ale with mint. A, B..., a, b... — means sharing the same letter in column are not significantly different ( $p < 0.05$ )

stances, including polyphenols, which diffuse into the beer during brewing and enhance its antioxidant capacity. This phenomenon is often accompanied by a decrease in pH and alcohol content, which may contribute to improved shelf life, stability, and potential health benefits (Borsa et al., 2022).

However, it should be emphasized that a high polyphenol content is not always beneficial. Excessive amounts may cause technological issues, such as turbidity or colloidal precipitation, as well as sensory defects – harsh or astringent taste notes (Humia et al., 2019). In lager beers, where a mild sensory profile and high clarity are expected, this is undesirable. Conversely, in PA, characterized by pronounced hop bitterness, a higher polyphenol level enhances the typical flavor profile and is considered desirable (Habschied et al., 2021). Therefore, optimizing polyphenol levels in beers enriched with herbal additives should balance health-promoting effects and sensory quality, which poses a significant challenge for brewing technologists.

Spearman rank correlation analysis revealed significant relationships between the studied parameters (Figure 3). The strongest correlations were observed between flavonoid content, TPC, and FRAP reducing capacity ( $r = 0.9-1.0$ ), clearly indicating that phenolic compounds – particularly flavonoids – are the main contributors to beer's

antioxidant activity. The ABTS<sup>+</sup> method also showed strong positive correlations with flavonoids, TPC, and FRAP ( $r = 0.9, 0.96, \text{ and } 0.9$ , respectively), confirming the consistency of results and the dominant role of these compounds in shaping beer's biological potential.

A negative correlation between pH and phenolic compound content and antioxidant activity ( $r = -0.59 \text{ to } -0.65$ ) was observed. Lower pH may promote increased extraction and stability of phenolic compounds from malt. Other studies (Chethan et al., 2007) indicate that mildly acidic beer conditions (pH 4–4.5) can favor the extraction process by improving polyphenol solubility, while higher pH conditions may decrease polyphenol content due to precipitation. It is also possible that the presence of phenolic compounds during wort fermentation may themselves contribute to pH reduction. Further studies are needed to clarify the exact mechanisms underlying these observations.

In contrast, beer color showed a positive correlation with flavonoid content and FRAP activity ( $r = 0.46$ ), implying that darker beers, enriched in Maillard compounds and polyphenols, tend to exhibit higher biological activity. The lack of significant correlation between alcohol content and antioxidant parameters indicates that ethanol itself is not responsible for beer's health-promoting properties, but




Figure 3. Spearman rank correlation matrix between antioxidant activity (ABTS, FRAP), phenolic compounds (flavonoids, total polyphenols), physicochemical parameters (pH, alcohol, real extract), and color (EBC). Significant positive correlations were observed between radical cation ABTS<sup>+</sup>, FRAP, flavonoids and total polyphenols, while pH showed a negative correlation with antioxidant parameters

rather the polyphenol fraction. Importantly, the positive correlation between extract, polyphenols, and antioxidant activity ( $r = 0.41\text{--}0.44$ ) suggests that the use of phenolic-rich raw materials and the extent of extraction are key factors influencing the antioxidant potential of the final product. Here, “antioxidant potential” is used to describe the beer’s capacity to neutralize free radicals, which is a more precise and measurable aspect of its biological activity. The extraction process applied in this study can be considered typical for the brewing conditions used; no comparative experiments with more intensive or alternative extraction methods were performed.

The obtained results clearly demonstrate that the content of phenolic compounds and flavonoids in beer is the main factor determining its antioxidant activity and indirectly influencing sensory characteristics such as color.

### Sensory evaluation

The results of the sensory evaluation of the PA beer (control sample) and its variants with herbal additives (thyme, ginger, lavender, marjoram and mint) are present-

ed in Figure 4 and Table 3. These outcomes clearly indicate that the adjuncts used significantly differentiated the flavor and aroma profiles of the beers, which was reflected both in the qualitative sensory characteristics and in consumer desirability ratings.

The highest desirability rating (Figure 4) received beer with mint (4.5 points), which was characterized by a fresh, cooling and slightly sweet taste and intense mint aroma. This profile was positively perceived by the panelists, which is also confirmed in the literature. Menthol and essential oils from mint are often suggested as factors enhancing the feeling of refreshment and improving the sensory acceptance of beverages (Hadi et al., 2025). Mint (*Mentha* spp.) belongs to *Lamiaceae* family and is rich in essential oils, polyphenols and flavonoids which a broad spectrum of biological activity (Best, 2022; Arshad et al., 2023) which is why it is frequently used as an additive in various products (Arshad et al., 2023).

The beer with ginger also achieved a high note (4.0 points) (Figure 4). It stood out with its spicy, aromatic aroma and slightly sweet flavor (Table 3). This sample was sensory similar to the base beer (control), indicating that




Figure 4. Sensory evaluation of examined beers: PA – Pale Ale (control), PAT – Pale Ale with thyme, PAG – Pale Ale with ginger, PAL – Pale Ale with lavender, PAMR – Pale Ale with marjoram, PAM – Pale Ale with mint. Rating scale: 1 – lowest desirability, 5 – highest desirability

Table 3. Results of sensory evaluation of examined beers.

Beer variant <sup>#</sup>	Flavor	Aroma	Color	Foam (density/color)
PA	Bitter, slightly hoppy	Hoppy, malty	Pale golden	Thick, white
PAT	Astringent, herbal, slightly bitter	Distinctly herbal, thyme	Pale amber	Medium, creamy
PAG	Spicy, slightly sweet, herbal	Spicy, fresh ginger	Golden	Fine, white
PAL	Bitter, slightly floral	Floral, lavender	Pale golden	Medium, whitish
PAMR	Herbal, slightly astringent	Herbal, dried herbs	Honey-like	Low, creamy
PAM	Fresh, cooling, slightly sweet	Minty, intense	Pale golden	Thick, white

<sup>#</sup> PA – Pale Ale (control), PAT – Pale Ale with thyme, PAG – Pale Ale with ginger, PAL – Pale Ale with lavender, PAMR – Pale Ale with marjoram, PAM – Pale Ale with mint

ginger blends well with the base profile of PA, without disrupting the beverage’s sensory balance. Ginger (*Zingiber officinale*) is a valuable botanical raw material, valued both for its flavor and health-promoting properties (Tozetto et al., 2019). It contains many bioactive compounds, including gingerols and shogaols, which are responsible for its intense aroma and functional properties (Haniadka et al., 2013). The literature highlights its anticancer activity (Zadorozhna, Mangieri, 2021; Zhao et al., 2020), antibacterial (Sebiomo et al., 2011; Srinivasan, 2017) and anti-inflammatory properties. The addition of ginger rhizome to the beer, not only enriches its sensory profile and imparts a warming effect, but may also enhance health benefits and microbiological stability (Sebiomo et al., 2011; Mao et al., 2019).

The lowest desirability was observed in beer with lavender, which received only 2.0 points (Figure 4). These samples were described as bitter and slightly floral (Table 3). This flavor and aroma profile appeared to be the least desirable among evaluators. Lavender flowers (*Lavandula angustifolia*) are known for their sedative and digestive properties (Matlawska, 2008) and possess an intense essential oil aroma containing over 100 volatile compounds (Kim et al., 2017; Wells et al., 2018). As a beer additive, it imparts a floral and bitter aroma (Table 3), described as spicy, floral, with notes of lemon and mint, giving the beer a unique flavor profile, and is sometimes used as a hop substitute in hop-free beers. Aromatic compounds from lavender may be perceived as dominant and discordant. These results align with FKelemen et al. (2024), who reported that Blonde Ale beers with lavender addition exhibited a strong lavender aroma during boiling, dominating the beer’s aroma, and increased bitterness (24.6 IBU vs. 15 IBU in control beers).

The different results were obtained for beers with the addition of thyme and marjoram (Figure 4). Both variants had a distinct herbal, astringent flavor, which resulted in moderate desirability among the evaluators (3.0–3.5 points). Herbal and slightly bitter notes were not perceived negatively, but did not increase the attractiveness of the beer, to the same extent as ginger or mint. A similar effect was observed by Djordjevic et al. (2016) during their studies on lager beers with herbal additives (lemon balm, thyme, juniper berries, hop cones), where they found that although thyme increased TPC, it did not have a positive effect on the sensory values of beer. Gościńska et al. (2019) and Solgajova et al. (2024) confirmed that thyme is not the best received beer addition. The beer with lemon balm and thyme addition was described by consumers as too herbal, resembling a „medicinal syrup”. The overall desirability and acceptability of product is determined by its flavor, aroma and color (Baryłko-Pikielna, Matuszewska, 2014). Flavor is the most important factor shaping desirability – as already discussed, the highest ratings were given to the beers with mint and ginger, while the lavender addition reduced the attractiveness of the product.

Similar effects were observed for aroma (Figure 4). The control beer had a characteristic hoppy and malty aroma (Table 3), typical of Pale Ale, and was rated as neutral (3.5 points). Beer with mint received the highest aroma score (4.5 points), described as minty and intense (Table 3). High marks were also given to beer with ginger (4.0 points), exhibiting subtle spicy notes (Table 3). Thus, mint and ginger enhanced aroma perception compared to the base beer. Other adjuncts received moderate aroma ratings: thyme and marjoram (3.0 points) were herbal and slightly astringent, while beer with lavender had the lowest aroma rating (2.0 points), characterized by an intense floral aroma.

Beer color was rated highly for all samples (Figure 4), indicating visual attractiveness. The highest scores (5.0 points) were given to beers with mint, ginger, and the base beer, described as pale gold, clear, and typical (Table 3). Beers with thyme and marjoram received slightly lower ratings (4.0 points), related to their darker color (pale amber, honey) (Table 3), but still highly acceptable. Color is a key selection indicator, being the first feature noticed by consumers. Market beers range from pale straw to golden, amber, brown, and black. In addition to Maillard reactions during malting and brewing, other reactions occur: caramelization, polyphenol oxidation, and pyrolysis, forming colored compounds. Polyphenol oxidation can occur during storage, particularly in light beers, leading to some darkening over time; however, this does not imply that polyphenol content increases due to oxidation. Darker beers generally contain higher initial levels of polyphenols, which strongly correlate with EBC, TPC, and flavonoid values.

The important attribute of beers is also the foam appearance (Bamforth, 2023). In conducted evaluations (Figure 4) all samples received high scores, from 4.0 points for the beer with marjoram, in which the foam was described as low and creamy (Table 3) to 5.0 in base beer and the mint variant, both characterized by dense and white foam (Table 3). Foam formation and stability in beer depend on the temperature of the beverage and its carbon dioxide content. The main substances responsible for foam formation are ethanol and lipids, while its endogenous stabilizers are polypeptides and bitter acids, further supported by melanoidins and metal ions. Therefore, the studied phenolic compounds play a minor role in this process (Bamforth, 2023). The observed differences in foam quality among the samples may result from interactions between proteins and polysaccharides and compounds derived from the herbal additives.

Principal component analysis (PCA) (Figure 5) visualized sensory differentiation among beer variants. PC1 accounted for 82.7% of total variance, and PC2 explained an additional 12.1%, indicating that these two axes capture nearly all variability. Clustering was evident: beers with thyme and marjoram were close, confirming similar sensory profiles (intense herbal notes, moderate desirability). The control beer and beer with ginger were positioned to-




Figure 5. Principal component analysis (PCA) in sensory evaluation of examined beers.

gether, indicating ginger did not significantly change the overall profile. Beer with mint occupied a distinct position, reflecting its unique, fresh, and highly rated profile. Lavender beer was separated on PC1, indicating its least accepted sensory profile. These results indicate that botanical additives differentiate sensory profiles and may also enhance functional properties through antioxidant compounds (polyphenols) and essential oils (Habschied et al., 2021; Galovicova et al., 2021; Silva et al., 2021). Thus, these raw materials are valuable for creating innovative beers, both in terms of sensory and health-promoting properties.

Beers with herbal adjuncts are attractive for both health-promoting properties and sensory qualities. The growing craft beer trend enables producers to experiment with formulas and reach new consumers. Herbal adjuncts, fruits, spices, and bee products allow for the creation of unique, innovative products targeting different groups.

Botanical ingredients contain a wide range of chemical constituents (polyphenols, vitamins, minerals) with health-promoting properties, enhancing sensory profile, stability, and shelf life (Solgajova et al., 2024; Coronas et al., 2024). However, careful process control is essential, as natural polyphenol degradation occurs during mashing, boiling, fermentation, and storage. Monitoring is also important in bottom-fermented beers (lagers), where high bioactive compound concentrations can cause cloudiness and sharp, intense aromas (Di Domenico et al., 2020; Habschied et al., 2021). Polyphenols have a dual role: they shape flavor (bitterness/astringency) and contribute to health benefits.

## CONCLUSIONS

The analysis conducted demonstrated that herbal adjuncts significantly differentiate the sensory profile of beer and substantially influence its consumer desirability. The highest ratings were given to beers with mint and ginger additions, while the lowest ratings were observed in beers containing lavender.

The PCA results confirmed clear clustering of sensory variants, indicating the possibility of categorizing herbal beers into more or less attractive groups for consumers. Botanical raw materials, in addition to shaping sensory characteristics, are a valuable source of bioactive compounds, such as polyphenols, which determine the beer's antioxidant properties. In summary, the use of botanical ingredients in craft beer production offers the opportunity to create unique and innovative products with enhanced health-promoting benefits and attractive sensory attributes.

## REFERENCES

- Ambra R., Pastore G., Lucchetti S., 2021.** The Role of Bioactive Phenolic Compounds on the Impact of Beer on Health. *Molecules*, 26, 486, <https://doi.org/10.3390/molecules26020486>.
- Anderson H.E., Santos I.C., Hildenbrand Z.L., Schug K.A., 2019.** A review of the analytical methods used for beer ingredient and finished product analysis and quality control. *Analytica Chimica Acta*, 1085(28): 1-20, <https://doi.org/10.1016/j.aca.2019.07.061>.
- Arshad M.K., Fatima I., Ahmad W., El-lahi S., Mumtaz M., Akhtar M.U., Siddique W.A., 2023.** Mint (*Mentha*): A herb and used as a functional ingredient. *Scholars International Journal of Traditional and Complementary Medicine*, 6(03): 38-52, doi: 10.36348/sijtc.2023.v06i03.003.
- Bamforth C.W., 2023.** The physics and chemistry of beer foam: A review. *European Food Research and Technology*, 249(1): 3-11, <https://doi.org/10.1007/s00217-022-04134-4>.
- Barylko-Pikielna N., Matuszewska I., 2014.** Sensoryczne Badania żywności. Podstawy. Metody. Zastosowania (Wyd. II). Kraków, Wydawnictwo Naukowe PTTŻ.
- Basařová G., Psota V., Šavel J., Basař P., Paulů R., Kosař K., Dostálék P., Basařová P., Kellner V., Mikulíková R., Čejka P., 2015.** Sladařství. Teorie a praxe výroby sladu (1st edition). Praha, Havlíček Brain Team, ISBN: 978-80-87109-47-2.
- Benzie I.F.F., Strain J.J., 1996.** The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the

FRAP assay. *Analytical Biochemistry*, 239: 70-76, <https://doi.org/10.1006/abio.1996.029224>.

- Best R., 2022.** Mint and menthol: A review of potential health benefits and wider human effects. doi: 10.20944/preprints202212.0273.v1.
- Borša A., Muntean M.V., Salanță L.C., Tofană M., Socaci S.A., Mudura, E., Pop A., Pop C.R., 2022.** Effects of Botanical Ingredients Addition on the Bioactive Compounds and Quality of Non-Alcoholic and Craft Beer. *Plants*, 11(15), 1958, <https://doi.org/10.3390/plants11151958>.
- Březinová M., 2021.** Beer Industry in the Czech Republic: Reasons for Founding a Craft Brewery. *Sustainability*, 13(17), 9680, <https://doi.org/10.3390/su13179680>.
- Carvalho D.O., Gonçalves L.M., Guido L.F., 2016.** Overall antioxidant properties of malt and how they are influenced by the individual constituents of barley and the malting process. *Comprehensive Reviews in Food Science and Food Safety*, 15: 927-943, doi: 10.1111/1541-4337.12218.
- Chethan S., Malleshi N.G., 2007.** Finger millet polyphenols: Optimization of extraction and the effect of pH on their stability. *Food chemistry*, 105(2): 862-870.
- Cocuzza S., Zarnkow M., Stallforth A., Peifer F., Jacob F., 2019.** The impact of dry hopping on selected physical and chemical attributes of beer. *BrewingScience*, doi: 10.23763/BrSc19-13cocuzza.
- Coronas R., Bianco A., Niccolai M., Fancello F., Sanna A.M.L., Asteggiano A., Medana C., Caboni P., Budroni M., Zara G., 2024.** Polyphenolic Content and Antimicrobial Effects of Plant Extracts as Adjuncts for Craft Herbal Beer Stabilization. *Foods*, 13(17): 2804, <https://doi.org/10.3390/foods13172804>.
- De Falco E., Mancini E., Roscigno G., Mignola E., Tagliata-tela-Scafati O., Senatore F., 2013.** Chemical Composition and Biological Activity of Essential Oils of *Origanum vulgare* L. subsp. *vulgare* L. under Different Growth Conditions. *Molecules*, 18(12): 14948-14960, <https://doi.org/10.3390/molecules181214948>.
- Di Domenico M., Feola A., Ambrosio P., et al., 2020.** Antioxidant Effect of Beer Polyphenols and Their Bioavailability in Dental-Derived Stem Cells (D-dSCs) and Human Intestinal Epithelial Lines (Caco-2) Cells. *Stem Cells International*, 2020: 8835813, doi: 10.1155/2020/8835813.
- Ditrych M., Kordialik-Bogacka E., Czyżowska A., 2016.** Antiradical and reducing potential of commercial beers. *Czech Journal of Food Sciences*, 33(3): 261-266.
- Djordjevic S., Popovic D., Despotovic S., Veljovic M., Atanackovic M., Cvejic J., Nedovic V., Leskosek-Cukalovic I., 2016.** Extracts of medicinal plants – As functional beer additives. *Chemical Industry and Chemical Engineering Quarterly*, 22: 301-308, doi: 10.2298/CICEQ150501044D.
- EBC-Analytica, EBC Analysis committee, 2010. 5th ed. Section 9 Beer. Method 9.43.2 Specific Gravity of Beer using a Density Meter; 9.4 Original, Real and Apparent Extract and Original Gravity of Beer; 9.8 Bitterness of Beer, Fachverlag.
- Elzinga K.G., Tremblay C.H., Tremblay V.J., 2015.** Craft beer in the United States: history, numbers, and geography. *Journal of Wine Economics*, 10 (3): 242-274, doi: 10.1017/jwe.2015.22.
- Fang H.H.P., Liang D.W., Zhang T., Liu Y., 2006.** Anaerobic treatment of phenol in wastewater under thermophilic

condition. *Water research*, 40 (3): 427-434, <https://doi.org/10.1016/j.watres.2005.11.025>.

- Galovičová L., Borotová P., Valková V., Vukovic N.L., Vukic M., Terentjeva M., Štefániková J., Ďuranová H., Kowalczewski P.L., Kačániová M., 2021.** *Thymus serpyllum* Essential Oil and Its Biological Activity as a Modern Food Preserver. *Plants*, 10(7), 1416, <https://doi.org/10.3390/plants10071416>.
- Gómez-Corona C., Escalona-Buendía H.B., García M., Chollet S., Valentin D., 2016.** Craft vs. industrial: Habits, attitudes and motivations towards beer consumption in Mexico. *Appetite*, 96: 358-367, doi: 10.1016/j.appet.2015.10.002.
- Gorjanovic S.Ž., Novakovic M.M., Potkonjak N.I., Ida L.C., Sužniewic D.Ž., 2010.** Application of a novel antioxidative assay in beer analysis and brewing process monitoring. *Journal of Agricultural and Food Chemistry*, 58: 744-751, doi: 10.1021/jf903091n.
- Gościnną K., Pobereźny J., Wszelaczyńska E., Łaba S., Świerczyński K., 2019.** Projektowanie funkcjonalnych piw rzemieślniczych o właściwościach prozdrowotnych, Uniwersytet Technologiczno-Przyrodniczy w Bydgoszczy, doi: 10.15199/65.2019.8.10.
- Gribkova I.N., Kharlamova L.N., Lazareva I.V., Zakharov M.A., Zakharova V.A., Kozlov V.I., 2022.** The influence of hop phenolic compounds on dry hopping beer quality. *Molecules*, 27(3), 740, <https://doi.org/10.3390/molecules27030740>.
- Guido L.F., Ferreira I.M., 2023.** The role of malt on beer flavour stability. *Fermentation*, 9(5), 464, <https://doi.org/10.3390/fermentation9050464>.
- Habschied K., Košir I.J., Krstanović V., Kumrić G., Mastanjević K., 2021.** Beer Polyphenols – Bitterness, Astringency, and Off-Flavors. *Beverages*, 7, 38, <https://doi.org/10.3390/beverages7020038>.
- Hadi C.S., Abbas K.S., Mohammed R.K., Hamasadeq Z.O., Hussein M.S., Hamad H.K., Hama S.A., 2025.** Development and characterization of mint powder (*Mentha spicata* L.)-fortified functional yoghurt: Physicochemical, sensory properties, and shelf-life assessment. *KUJAS*, 16, 2: 176-183, doi: 10.58928/ku24.16222.
- Haniadka R., Saldanha E., Sunita V., Palatty P.L., Fayad R., Baliga M.S., 2013.** A review of the gastroprotective effects of ginger (*Zingiber officinale* Roscoe). *Food & Function*, 4(6): 845-855, doi: 10.1039/c3fo30337c.
- Horincar G., Enachi E., Bolea C., Râpeanu G., Aprodu I., 2020.** Value-Added Lager Beer Enriched with Eggplant (*Solanum melongena* L.) Peel Extract. *Molecules*, 25, #731, <https://doi.org/10.3390/molecules25030731>.
- Humia B.V., Santos K.S., Barbosa A.M., Sawata M., Mendonça M.d.C., Padilha F.F., 2019.** Beer molecules and its sensory and biological properties: A review. *Molecules*, 24, 1568, <https://doi.org/10.3390/molecules24081568>.
- Kelemen M., Albert Cs., András Cs.D., Salamon R.V., 2024.** Preliminary study of lavender-flavoured beer production methods. *Acta Univ. Sapientiae, Alimentaria*, 17: 120-135, doi: 10.47745/ausal-2024-0007.
- Kikut-Ligaj D., Mikołajczyk-Bator K., 2022.** Analiza preferencji konsumentów wobec piw oraz sterowanie smakiem gorzkim piw chmielonych. *Engineering Sciences and Technologies*, 1(38): 90-110, doi: 10.15611/nit.2022.38.06.

- Kim Ch.Y., Jang K.S., Kwon O.H., Jeon S.G., Kwon J.B., Dhungana S.K., Mun J.H., Park Y.S., Kim I.D., 2017.** Addition of green pepper enhanced antioxidant potential and overall acceptance of beer. *International Journal of Sciences*, 6(05): 49-54, doi: 10.18483/ijSci.1288.
- Kumari B., Sit N., 2025.** Filtration and clarification. pp. 85-120. In: *Mechanical Separation Processes in the Food Industry*. Woodhead Publishing, DOI:10.1016/B978-0-12-819523-9.00008-1.
- Lehrl R., 2017.** Domowe warzenie piwa (wyd. 6). Warszawa, Wydawnictwo RM.
- Lin R., Chen X., Peng Z. et al., 2023.** Evaluation of polyphenols content and antioxidant ability of commercial herbal beer. *Systems Microbiology and Biomanufacturing*, 4: 77-85, <https://doi.org/10.1007/s43393-023-00185-0>.
- Mao Q.Q., Xu X.Y., Cao S.Y., Gan R.Y., Corke H., Beta T., Li H.B., 2019.** Bioactive Compounds and Bioactivities of Ginger (*Zingiber officinale Roscoe*). *Foods*, 8(6), #185, doi: 10.3390/foods8060185.
- Mastanjević K., Krstanović V., Lukinac J., Jukić M., Lucan M., Mastanjević K., 2019.** Craft brewing – is it really about the sensory revolution? *Kvasny prumysl*, 65: 13-16, <https://doi.org/10.18832/kp2019.65.13>.
- Matlawska I., (2008).** Farmakognozja. UAM Poznań.
- Mazengia G., Dessalegn E., Dessalegn T., 2022.** Effect of *Moringa stenopetala* leaf extracts on the physicochemical characteristics and sensory properties of lagered beer. *Food Science & Nutrition*, 10: 507-514, <https://doi.org/10.1002/fsn3.2672>.
- Nardini M., Garaguso I., 2020.** Characterization of Bioactive Compounds and Antioxidant Activity of Fruit Beers. *Food Chemistry*, 305, #125437, PMID: 31499290; 10.1016/j.foodchem.2019.125437.
- Olšovská J., Matoulková D., Čejka J., Jurková M., 2014.** Beer and Health. *Kvasný průmysl*. 60(7-8): 174-181, <https://doi.org/10.18832/kp2014017>.
- Papoutsis K., Pristijono P., Golding J.B., Stathopoulos C.E., Bowyer M.C., Scarlett C.J., Vuong Q.V., 2016.** Optimisation of aqueous extraction conditions for the recovery of phenolic compounds and antioxidants from lemon pomace. *International Journal of Food Science & Technology*, 51(9): 2009-2018, <https://dx.doi.org/10.1111/ijfs.13168>.
- Pieczonka S.A., Hemmler D., Moritz F., Lucio M., Zarnkow M., Jacob F., Rychlik M., Schmitt-Kopplin P., 2021.** Hidden in its Color: A molecular-level analysis of the beer's Maillard reaction network. *Food Chemistry*, vol. 361, DOI: 10.1016/j.foodchem.2021.130112.
- Pluháčková H., Gregor T., Boško R., Běláková S., Svoboda Z., Benešová K., 2020.** Fortification of Beer with Extracts of the Selected Czech Medicinal Herbs and Plants. *Kvasný průmysl*, 66(4): 314-319, <https://doi.org/10.18832/kp2019.66.314>.
- Polska Norma PN-90/A-75101/06 Przetwory owocowe i warzywne. Przygotowanie próbek i metody badań fizykochemicznych. Oznaczenie pH metodą potencjometryczną.
- Preiss R., Tyrawa C., Krogerus K., Garshol L.M., van der Merwe G., 2018.** Traditional Norwegian kveik are a genetically distinct group of domesticated *Saccharomyces cerevisiae* brewing yeasts. *Frontiers in Microbiology*, 9, #2137, <https://doi.org/10.3389/fmicb.2018.02137>.
- Ramos da Silva L.R., Ferreira O.O., Cruz J.N., Pereira Franco C.J., Dos Anjos T.O., Cascaes M.M., Almeida da Costa W., Andrade E.H.A., 2021.** *Lamiaceae* Essential Oils, Phytochemical Profile, Antioxidant, and Biological Activities. Evidence-based Complementary and Alternative Medicine, pp. 1-18., <https://doi.org/10.1155/2021/6748052>
- Raport UOKiK <https://uokik.gov.pl/raport-uokik-polski-rynek-piwa-i-chmielu> [accessed 17.06.2025]
- Re R., Pellegrini N., Proteggente A., Pannala A., Yang M., Rice-Evans C., 1999.** Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, 26(9-10): 1231-1237, doi: 10.1016/s0891-5849(98)00315-3.
- Rum L., 2005.** Ilustrowany leksykon piwa. Wydawnictwo Kurpisz, ISBN: 978-83-75249-84-2.
- Sanna V., Pretti L., 2015.** Effect of wine barrel ageing or *sapa* addition on total polyphenol content and antioxidant activities of some Italian craft beers. *International Journal of Food Science + Technology*, 50: 700-707, <https://doi.org/10.1111/ijfs.12666>.
- Sebiomo A., Awofodu A.D., Awosanya A.O., Awotona F.E., Ajayi A.J., 2011.** Comparative studies of antibacterial effect of some antibiotics and ginger (*Zingiber officinale*) on two pathogenic bacteria. *Journal of Microbiology and Antimicrobials*, 3(1): 18-22.
- Shopska V., Denkova-Kostova R., Dzhivoderova-Zarcheva M., Teneva D., Denev P., Kostov G., 2021.** Comparative study on phenolic content and antioxidant activity of different malt types. *Antioxidants*, 10(7): 1124, <https://doi.org/10.3390/antiox10071124>.
- Silva S., Cruz A., Oliveira R.F., Oliveira A.I., Pinho C., 2021.** Potential biological activities of craft beer: A review. *Acta Portuguesa de Nutrição*, 25: 84-89, doi: 10.21011/apn.2021.2515.
- Skendi A., Irakli M., Chatzopoulou P., 2017.** Analysis of phenolic compounds in Greek plants of *Lamiaceae* family by HPLC. *Journal of Applied Research on Medicinal and Aromatic Plants*, 6: 62-69, <https://doi.org/10.1016/j.jar-map.2017.02.001>.
- Sohrabvandi S., Mortazavian A., Rezaei K., 2012.** Health-related aspects of beer: a review. *International Journal Food Properties*, 15(2): 350-373, <http://doi.org/10.1080/10942912.2010.487627>.
- Sokol-Łętowska A., 2013.** Związki fenolowe w nalewkach z wybranych owoców. UWP Wrocław.
- Solgajová M., Dráb Š., Straka D., Mendelová A., Kolesárová A., Mareček J., 2024.** Effect of the addition of selected herbs on the technological and sensory quality of beer. *Journal of microbiology, biotechnology and food sciences*, e11012, doi: 10.55251/jmbfs.11012.
- Sperkowska B.M., Wojtynek C., Walkowiak S., Koba M., 2021.** Porównanie potencjału antyoksydacyjnego i zawartości składników fenolowych w piwach konwencjonalnych i bezalkoholowych. *Journal of Health & Environmental Research/Bromatologia & Chemia Toksykologiczna*, 54(3), doi: 10.32383/bct/157163.
- Srinivasan K., 2017.** Ginger rhizomes (*Zingiber officinale*): A spice with multiple health beneficial potentials. *PharmaNutrition*, 5(1): 18-28, <https://doi.org/10.1016/j.phanu.2017.01.001>.
- Šibalić D., Planinić M., Jurić A., Bucić-Kojić A., Tišma M., 2021.** Analysis of phenolic compounds in beer: from raw ma-

- terials to the final product. *Chemical Papers*, 75(1): 67-76, doi: 10.1007/s11696-020-01276-1.
- Tozetto L.M., do Nascimento R.F., de Oliveira M.H., Van Beik J., Canteri M.H.G., 2019.** Production and physicochemical characterization of craft beer with ginger (*Zingiber officinale*). *Food Science & Technology*, 39(4): 962-970, <https://doi.org/10.1590/fst.16518>.
- Ullah A., Munir S., Badshah S.L., Khan N., Ghani L., Poulson B.G., Emwas A.H., Jaremko M., 2020.** Important flavonoids and their role as a therapeutic agent. *Molecules*, 25(2), 5243, <http://doi.org/10.3390/molecules25225243>.
- Ualema N.J.M., Dos Santos L.N., Bogusz S., Ferreira N.R., 2024.** From Conventional to Craft Beer: Perception, Source, and Production of Beer Color-A Systematic Review and Bibliometric Analysis. *Foods*, 13(18): 2956, doi: 10.3390/foods13182956.
- Viejo C.G., Fuentes S., Torrico D., Howell K., Dunshea F.R., 2018.** Assessment of beer quality based on foamability and chemical composition using computer vision algorithms, near infrared spectroscopy and machine learning algorithms. *Journal of Science of Food and Agriculture*, 98(2): 618-627, <https://doi.org/10.1002/jsfa.8506>.
- Wells R., Truong F., Adal A.M., Sarker L.S., Mahmoud S.S., 2018.** Lavandula essential oils: A Current Review of Applications in Medicinal, Food, and Cosmetic Industries of Lavender. *Natural Product Communications*, 13(10): 1403-1417, doi: 10.1177/1934578X1801301038.
- Wojtyra B., 2020.** How and why did craft breweries „revolutionise” the beer market? The case of Poland. *Moravian Geographical Reports*, 28: 81-97, <https://doi.org/10.2478/mgr-2020-0007>.
- Yalçınçıray Ö., Vural N., Anlı R.E., 2022.** Effects of filtration and pasteurization process on bioactive phenolic compounds of beer. *Journal of Food Processing and Preservation*, 46(2), <https://doi.org/10.1111/jfpp.16234>.
- Zadorozhna M., Mangieri D., 2021.** Mechanisms of chemopreventive and therapeutic properties of ginger extracts in cancer. *International Journal of Molecular Sciences*, 22(12), #6599, doi: 10.3390/ijms22126599.
- Zhao L., Rupji M., Choudhary I., Osan R., Kapoor S., Zhang H.J., Yang C., Aneja R., 2020.** Efficacy based ginger fingerprinting reveals potential antiproliferative analytes for triple negative breast cancer. *Scientific Reports*, 10(1), #19182, doi: 10.1038/s41598-020-75707-0.